# <u>Congruence Of Triangles And Inequalities in a Triangle</u> CHAPTER 9

Exercise – 9 (A)

Answer1) Given: AB || CD D **<u>To prove</u>**: i) 0 is the midpoint of AD ii)  $\triangle AOB \cong \triangle DOC$ **Proof:** In  $\triangle AOB$  and  $\triangle DOC$ OA = OD(Given) (vertically opposite  $\angle AOB = \angle COD$ ametextbooks, W В angles)  $\angle OAB = \angle ODC$ (alternate angles) Therefore; (A.A.S. criteria)  $\Delta AOB \cong \Delta DOC$ Hence; (c.p.c.t.) OB = OCHence proved. Answer2) в С Given: AD=BD To prove: CD bisects AB i.e. OA=OB Ó Proof: In  $\triangle$ BOC and  $\triangle$ AOD

**CLASS IX** 



 $\angle OAD = \angle OBC = 90$ (Given)

∠AOD=∠BOC (vertically opp. Angles)

(A..A.S criteria) Therefore  $\triangle BOC \cong \triangle AOD$ 

Hence OA=OB i.e CD bisects AB

Hence proved.

# Answer3)

<u>Given</u>: (i) l || m

(ii) p || q

**<u>To prove</u>**:  $\triangle ABC \cong \triangle CDA$ 

# Proof:

verversal ... angles) 4 (Alternate angles) common) Taking | parallel to m AC is the trAnswerversal

 $\angle ACB = \angle CAD$ 

Taking p parallel to q

∠BAC=∠DCA

AC=AC

Therefore;


 $\Delta ABC \cong \Delta CDA$ (ASA ctiteria)

Hence Proved.

Answer4)Given: AB=AC

**<u>To prove</u>**: (i)AD bisects BC (i.e. BD=DC)

(ii) AD bisects  $\angle A$ 



D


С

m

С



## Prove:



# Answer6)

# Given:

(i)  $\triangle$ ABC and  $\triangle$ DBC are two isosceles triangles in which AB=AC & BD=DC.

D

Е

# To Prove:

- (i)  $\triangle ABD \cong \triangle ACD$
- (ii)  $\triangle ABE \cong \triangle ACP$
- (iii) AE bisects  $\angle A$  as well as  $\angle D$ .
- (iv) AE is the perpendicular bisector of BC.

## Proof:

(i) In  $\triangle$ ABD and  $\triangle$ ACD, AD = AD (Common) AB = AC (Given). (SSS criteria) BD = CD (Given) Therefore,  $\triangle ABD \cong \triangle ACD$  $\angle BAD = \angle CAD(C.P.C.T)$  $\angle BAE = \angle CAE$ (ii) In  $\triangle ABE \& \triangle ACE$ AE = AE (Common)  $\angle BAE = \angle CAE$ (Proved above) AB = AC (Given) Therefore,

#### **CLASS IX**

```
ΔABE≅ ΔACE
                               (SAS criteria).
(iii) \angle BAD = \angle CAD (proved in part i)
Hence, AE bisects \angle A.
also,
In \triangleBED and \triangleCED
ED = ED (Common)
BD = CD (Given)
BE = CE
                                  A as ZD
(c.p.c.t)
(\Delta ABE \cong \Delta ACE \text{ so by c.p.c.t})
Therefore, \triangle BED \cong \triangle CED
Thus,
\angle BDE = \angle CDE
                        ( c.p.c.t)
Hence, we can say that AE bisects \angle A as well as \angle D.
(iv) \angle BED = \angle CED
(by CPCT as \triangle BED \cong \triangle CED)
Therefore;
BE = CE
\angle BED + \angle CED = 180^{\circ} (BC is a straight line)
\Rightarrow 2 \angle BPD = 180^{\circ}
\Rightarrow \angle BED = 90^{\circ}
```

Hence, AE is the perpendicular bisector of BC.

# **CLASS IX**

# Answer7)

Given: (i)x=y

(ii) AB=CB

To prove: AE = CD

Proof:

Consider the triangles AEB and CDB.

∠EBA=∠DBC∠EBA=∠DBC (Common angle) ...(i)

Further, we have:

∠BEA=180-y

∠BDC=180-x

Since x = y,

we have:

180-x = 180-y

⇒∠BEA=∠BDC ...(ii)

AB = CB (Given) ...(iii) From (i), (ii) and (iii),

we have:  $\triangle BDC \cong \triangle BEA$  (AAS criterion)  $\therefore AE = CD (CPCT)$ Hence, proved.

А

D

x٥

# Answer8)

**<u>Given</u>**: (i) l is the bisector of an  $\angle A$ 

(ii) BP and BQ are perpendiculars

**<u>To Prove</u>**: ΔAPB≅ΔAQB

Proof:

In  $\triangle APB$  and  $\triangle AQB$ 

 $\angle P = \angle Q$  (Right angles)

 $\angle BAP = \angle BAQ$  (l is the bisector)

(A.A.S criteria)

Retextbooks,

AB = AB (Common)

 $\Delta APB \cong \Delta AQB$ 

Hence, Proved.

(ii) BP = BQ (By c.p.c.t)


Therefore,

B is equidistant from the arms of  $\angle A$ 

# Answer9)

Given: AC bisects angles A and C.

 $\frac{\text{To prove}}{\text{(ii) } AB} = AD$ (ii) CB = CD



В



# Proof:

 $\Delta$  ABC and  $\Delta$ ADC,

we have:

 $\angle CAB = \angle CAD$ 

 $\angle$  BCA =  $\angle$  DCA

AC = AC (common)

 $\triangle ABC \cong \triangle ADC$ 

Therefore,

AD = AB(c.p.c.t)

CD = BC(c.p.c.t)

Answer10)Given: AB=AC

To prove: AC+AD=BC

Proof:

Let AB = AC = a and AD = b

In a right angled triangle ABC ,  $BC^2 = AB^2 + AC^2$ 

 $BC^2 = a^2 + a^2$ 

BC =  $a\sqrt{2}$ 

Given AD = b, we get

DB = AB - AD or DB = a - b



We have to prove that AC + AD = BC or  $(a + b) = a\sqrt{2}$ .

By the angle bisector theorem, we get

AD/ DB = AC / BC  

$$b/(a - b) = a/a\sqrt{2}$$

$$b/(a - b) = 1/\sqrt{2}$$

$$b = (a - b)/\sqrt{2}$$

$$b\sqrt{2} = a - b$$

$$b(1 + \sqrt{2}) = a$$

$$b = a/(1 + \sqrt{2})$$
Rationalizing the denominator with  $(1 - \sqrt{2})$ 

$$b = a(1 - \sqrt{2}) / (1 + \sqrt{2}) \times (1 - \sqrt{2})$$

$$b = a(1 - \sqrt{2}) / (-1)$$

$$b = a(\sqrt{2} - 1)$$

$$b = a\sqrt{2} - a$$

or AD + AC = BC [we know that AC = a, AD = b and  $BC = a\sqrt{2}$ ]

Hence it is proved.

# Answer11)

Given: ( i) 0A=0B

(ii)0P=0Q

To Prove: (i) PX=QX

(ii)AX=BX

# Proof:

In  $\Delta$  PBO and  $\Delta AOQ$ 

OB=OA (Given)

 $\begin{array}{c} OP=OQ \quad (Given) \\ \label{eq:generalized} & \\ \end{tabular} \\ C= \end{tabular} 0 \quad (Given) \\ \end{tabular} \\ Therefore; \\ \end{tabular} \\ \end{tabular}$ 

Answer12)

Given: (i) ABC is an equilateral triangle,

(ii) PQ  $\parallel AC$ 

(iii) CR=BP

**<u>To prove</u>**: QR bisects PC or PM = MC

Proof:

**CLASS IX** 

Since,  $\triangle ABC$  is equilateral triangle,

$$\angle A = \angle ACB = 60^{\circ}$$

Since, PQ ||AC and corresponding angles are equal,

 $\angle BPQ = \angle ACB = 60^{\circ}$ 

In ΔBPQ,

 $\angle B = \angle ACB = 60^{\circ}$ 

 $\angle BPQ = 60^{\circ}$ 

Hence,  $\Delta$ BPQ is an equilateral triangle.

 $\therefore$  PQ = BP = BQ

Since we have BP = CR,

We say that  $PQ = CR \dots (1)$ 

Consider the  $\Delta PMQ$  and  $\Delta CMR$ ,

 $\angle PQM = \angle CRM$ 

(alternate angles)

(AAS criteria)

 $\angle PMQ = \angle CMR$  (vertically opposite angles)

 $PQ = CR \dots from 1$ 

 $\Delta PMQ \cong \Delta CMR$ 

 $\therefore PM = MC \qquad (c.p.c.t)$ 


Hence proved.

# Answer13)

Given: (i)AB ll DC



HDOOMS, HINCH, BUILD



 $\underline{\text{To Prove}}$ : (i) AB = CQ (ii) DQ = DC + ABso, AB ll DQ so,  $\angle BAQ = \angle DQA$  (alternate angles) or  $\angle$  BAP =  $\angle$  CQP -----(1) Now, in triangle ABP and triangle QCP,  $\angle$  BAP =  $\angle$  CQP (from (1))  $\angle$  BPA =  $\angle$  CPQ (vertically opposite angles) BP = CP (since P is the midpoint of BC) so, triangle ABP congruent triangle QCP (by AAS congruency) or AB = CQ (by CPCT) [proved] -----(2) again, DQ = DC + CQ = DC + AB (from (2)) [proved] Answer14) **<u>Given</u>**: ABCD is a square and PB=PD To prove: CPA is a straight line

Proof:

 $\Delta$ APD and  $\Delta$ APB,

 $DA = AB \dots$ (as ABCD is square)

 $AP = AP \dots$ 

(common side)

$PB = PD \dots$	(Given)
$\Delta APD \cong \Delta APB$	(SSS criteria)

Hence, we know that, corresponding parts of the congruent triangles are equal

```
\angle APD = \angle APB \dots (1)
```

 $C^{*}$ 

# **CLASS IX**

Now consider  $\triangle$ CPD and  $\triangle$ CPB,

 $CD = CB \dots ABCD$  is square

 $CP = CP \dots$  common side

 $PB = PD \dots Given$ 

Thus by SSS property of congruence,

 $\Delta CPD \cong \Delta CPB$ 

 $\angle CPD = \angle CPB \dots (C.P.C.T.)\dots(2)$ 

Now,

Adding both sides of 1 and 2,

 $\angle CPD + \angle APD = \angle APB + \angle CPB \dots (3)$ 

Angels around the point P add upto 360°

 $\therefore \angle CPD + \angle APD + \angle APB + \angle CPB = 360^{\circ} \dots (4)$ 

From 4,

 $2(\angle CPD + \angle APD) = 360^{\circ}$  $\angle CPD + \angle APD = 180^{\circ}$ 

This proves that CPA is a straight line.


**Answer15**) <u>Given</u>: In square ABCD,  $\triangle$ OAB is an equilateral triangle.

**To prove:**  $\triangle OCD$  is an isosceles triangle.

#### **Proof:**

 $\therefore \angle DAB = \angle CBA = 90^{\circ}$ (Angles of square ABCD)

(Angles of equilateral  $\triangle OAB$ ) And,  $\angle OAB = OBA = 60^{\circ}$ 



# **CLASS IX**

 $\therefore \angle DAB - \angle OAB = \angle CBA - \angle OBA = 90^{\circ} - 60^{\circ}$ 

⇒∠OAD=∠OBC=30° ....(i)

∵∠DAB=∠CBA=90° Angles of square ABCD

And,  $\angle OAB = OBA = 60^{\circ}$ 

Angles of equilateral  $\Delta OAB$ 

∴∠DAB-∠OAB=∠CBA-∠OBA=90°-60°

 $\Rightarrow \angle OAD = \angle OBC = 30^{\circ}$ ....(i)

Now, in  $\Delta DAO$  and  $\Delta CBO$ ,

AD = BC(Sides of square ABCD)  $\angle DAO = \angle CBO$ [From (i)] (Sides of equilateral  $\triangle OAB$ ) AO = BO

∴ By SAS congruence criteria,  $\Delta DAO \cong \Delta CBO$ 

So, OD = OC(CPCT) Hence,  $\triangle OCD$  is an isosceles triangle.

Answer16)

<u>Given:</u> AX = AY.


To prove: CX = BY

Proof:

In  $\Delta CXA$  and  $\Delta BYA$ ,

AX = AY .....Given

 $\angle XAC = \angle YAB \dots$  common angle






 $AC = AB \dots Given$ ,

 $\Delta CXA \cong \Delta BYA$  (S.A.S. criteria)

CX = BY (C.P.C.T.)

Answer17)



Hisch awa

**<u>Given:</u>** BD = DC and  $DL \perp AB$  and  $DM \perp AC$  such that DL=DM

To prove: AB = AC

## Proof:

In right angled triangles  $\Delta$ BLD and  $\Delta$ CMD,

 $\angle BLD = \angle CMD = 90^{\circ}$ 

 $BD = CD \dots Given$ 

 $DL = DM \dots Given$ 

Thus by right angled hypotenuse side property of congruence,

 $\Delta BLD \cong \Delta CMD$ 

Hence, we know that, corresponding parts of the congruent triangles are equal

 $\angle ABD = \angle ACD$ 


In  $\triangle$ ABC, we have,

 $\angle ABD = \angle ACD$ 

 $\therefore$  AB= AC .... Sides opposite to equal angles are equal

# Answer18)

**<u>Given</u>**: In  $\triangle$ ABC, AB=AC and the bisectors of  $\angle$ B and  $\angle$ C meet at a point O.



**To prove:** BO=CO and  $\angle$ BAO =  $\angle$ CAO

# Proof:

In ,  $\triangle$  ABC we have,

$$\therefore \text{ OC} = \text{OB} \dots (1)$$

...ven  $\zeta = 20$  cm  $\zeta = 20$ 

 $\angle ABO = 2 \angle B$ 1  $\angle ACO = \frac{1}{2} \angle C$  $\therefore \angle ABO = \angle ACO \dots (2)$ Now in  $\triangle ABO$  and  $\triangle ACO$  $AB = AC \dots Given$ 

 $\angle ABO = \angle ACO \dots$  from (2)

BO = OC ... from (1)


Thus by SAS property of congruence,

 $\Delta ABO \cong \Delta ACO$ 

Hence, we know that, corresponding parts of the congruent triangles are equal

 $\angle BAO = \angle CAO$ 

ie. AO bisects  $\angle A$ ; Hence proved.



**CLASS IX** 

 $\angle$ MNA= $\angle$ BNM(CPCT) ....(1)

 $\angle$ MNC= $\angle$ MND=90 ....(2)

Subtracting Eq(2) from Eq(1)

∠MND-∠MNA=∠MNC-∠BNM

∠AND=∠BNC

 $\Delta$ AND congruent to  $\Delta$ BNC

AD = BC(CPCT)

Hence proved

Answer20)

<u>**Given:</u>** Bisectors of the angles B and C of an isosceles triangle with AB = AC intersect</u> ametextbool each other at O.

A BHB

в

М

<u>To prove:</u> $\angle$ MOC =  $\angle$ ABC

Proof:

In  $\triangle ABC$ ,

AB = AC (Given)

 $\Rightarrow \angle ACB = \angle ABC$  (opposite angles to equal sides are equal)

 $1/2 \angle ACB = 1/2 \angle ABC$  (divide both sides by 2)

 $\Rightarrow \angle OCB = \angle OBC \dots (1)$  (As OB and OC are bisector of  $\angle B$  and  $\angle C$ )

Now,  $\angle MOC = \angle OBC + \angle OCB$  (as exterior angle is equal to sum of two opposite interior angle)

 $\Rightarrow \angle MOC = \angle OBC + \angle OBC$  (from (1))

 $\Rightarrow \angle MOC = 2 \angle OBC$ 

 $\Rightarrow \angle MOC = \angle ABC$  (because OB is bisector of  $\angle B$ )

Hence proved.

**Answer21)** <u>Given:</u> (i) In an isosceles ΔABC,

(ii) AB = AC,

(iii) BO and CO are the bisectors of  $\angle$ ABC and  $\angle$ ACB.

To prove: ∠ABD = ∠BOC Construction: Produce CB to point D. Proof:

In ∆ABC,

 $\therefore AB = AC$ (Given)  $\therefore \angle ACB = \angle ABC$ (Angle opposite to equal sides are equal)

 $\Rightarrow \frac{1}{2} \angle ACB = \frac{1}{2} \angle ABC$ 

⇒∠0CB=∠0BC .....(i) (Given, B0 and C0 are angle bisector of ∠ABC and ∠ACB, respectively)


 $\Rightarrow \frac{1}{2} \angle ACB = \frac{1}{2} \angle ABC$ 

 $\Rightarrow \angle OCB = \angle OBC$  .....(i) (Given, BO and CO are angle bisector of  $\angle ABC$  and  $\angle ACB$ , respectively)

In  $\Delta BOC$ ,

 $\angle OBC + \angle OCB + \angle BOC = 180^{\circ}$  (By angle sum property of triangle)

 $\Rightarrow \angle OBC + \angle OBC + \angle BOC = 180^{\circ}$  [From (i)]



**CLASS IX** 

0r,

 $\angle BPQ = \angle PBQ$ 

Now, in  $\triangle$ BPQ

 $\angle BPQ = \angle PBQ$ 

 $\Delta BPQ$  is an isosceles triangle

Hence Proved.

Answer23) Given: A is an object in front of mirror LM,

B is the image of A and the observer is at D

AB intersects LM at T

To Prove: A and B are equ	aidistant from LM
AT = BT	A Har
Construction: Join BD. Let it intersect LM at C	
Join AC. CN be the normal at C.	
Proof:	ter.
$\angle i = \angle r$	(1)
AB   NC	[Both are perpendicular to LM]
$\angle CAT = \angle CAN = \angle i$	(2)[Alternate angles]
$\angle CBA = \angle DCN = \angle r$	(3)[Corresponding angles]

From (1), (2) and (3), we get

...(4)  $\angle CAT = \angle CBA$ 

In  $\triangle$ CAT and  $\triangle$ CBT,

 $\angle CAT = \angle CBT$ ...[From (4)]

**CLASS IX** 

**RS** Aggarwal solutions

Μ

D

N

$\angle ATC = \angle BTC$	[Each 90°]
CT = CT	[Common side]
Therefore;	
$\Delta CAT \cong \Delta CBT$	[ AAA Criteria]
AT = BT	[C.P.C.T]
Hence Proved.	
Answer24)	

Let AB be the breadth of the river. M is any point situated on the bank of the river. Let 0 be the mid point of BM.

Moving along perpendicular to point such that A , ,O and N are in straight line.

Then MN is the required breadth of the river. enthooksill

In  $\triangle OBA$  and  $\triangle OMN$ ,

we have:OB=OM (0 is midpoint)

 $\angle OBA = \angle OMN$  (Each 90°)

```
\angle AOB = \angle NOM (Vertically opposite angle)
```

 $\therefore \triangle OBA \cong \triangle OMN$  (ASA criterion)

In  $\triangle OBA$  and  $\triangle OMN$ .

we have:OB=OM (0 is midpoint)

 $\angle OBA = \angle OMN$  (Each 90°)

 $\angle AOB = \angle NOM$  (Vertically opposite angle)

 $\therefore \triangle OBA \cong \triangle OMN$  (ASA criterion) Thus, MN = AB (CPCT) If MN is known, one can measure the width of the river without actually crossing it.

**CLASS IX** 

**Answer25)**Given: D is the midpoint of ac  $BD = \frac{1}{2} AC$ 

To Prove: ∠ABC is 90°

In  $\triangle$ ADB, AD = BD

 $\angle DAB = \angle DBA = \angle x$ 

(Opposite angles)

In  $\Delta DCB$ , BD = CD

 $\angle DBC = \angle DCB = \angle y$ 

In  $\triangle$ ABC we will use the angle sum property

 $\angle ABC + \angle BCA + \angle CAB = 180^{\circ}$ 

- $2(\angle x + \angle y) = 180^{\circ}$
- $\angle x + \angle y = 90^{\circ}$


 $\angle ABC = 90^{\circ}$ 

This meAnswer that ABC is the right angled triangle.

Answer26)No, because in the congruent rule, the two sides and the included angle of one triangle are equal to the two sides and the included angle of the other triangle, that is, SAS criteria.

Answer 27) No,

Corresponding sides must be equal.



**CLASS IX** 

# EXERCISE-9(B)

**Answer1)** (i) No, because the sum of two sides of a triangle is not greater than the third side.

5 + 4 = 9

(ii) Yes, because the sum of two sides of a triangle is greater than the third side.

7 + 4 > 8; 8 + 7 > 4; 8 + 4 > 7

(iii) Yes, because the sum of two sides of a triangle is greater than the third side.

5 + 6 > 10; 10 + 6 > 5; 5 + 10 > 6

(iv) Yes, because the sum of two sides of a triangle is greater than the third side. 2.5 + 5 > 7; 5 + 7 > 2.5; 2.5 + 7 > 5

(v) No, because the sum of two sides of a triangle is not greater than the third side. 3 + 4 < 8

**Answer2**) <u>Given</u>: In  $\triangle ABC$ ,  $\angle A = 50^{\circ}$  and  $\angle B = 60^{\circ}$ 

In ΔABC,

 $\angle A + \angle B + \angle C = 180^{\circ}$  (Angle sum property of a triangle)

 $\Rightarrow$ 50° + 60° +  $\angle$ C = 180°

 $\Rightarrow 110^{\circ} + \angle C = 180^{\circ}$ 

 $\Rightarrow \angle C = 180^{\circ} - 110^{\circ}$  $\Rightarrow \angle C = 70^{\circ}$ 

Hence, the longest side will be opposite to the largest angle ( $\angle C = 70^\circ$ ) i.e. AB. And, the shortest side will be opposite to the smallest angle ( $\angle A = 50^\circ$ ) i.e. BC.

**Answer3)** (i) <u>Given</u>: In  $\triangle$ ABC,  $\angle$ A = 90°

So, sum of the other two angles in triangle  $\angle B + \angle C = 90^{\circ}$ 

i.e.  $\angle B$ ,  $\angle C < 90^{\circ}$ 

Since,  $\angle A$  is the greatest angle.

So, the longest side is BC.

(ii) Given:  $\angle A = \angle B = 45^{\circ}$ 

Using angle sum property of triangle,

 $\angle C = 90^{\circ}$ 

Since,  $\angle C$  is the greatest angle.

So, the longest side is AB.

(iii) Given:  $\angle A = 100^{\circ}$  and  $\angle C = 50^{\circ}$ 

Using angle sum property of triangle,

 $\angle B = 30^{\circ}$ 

Since,  $\angle A$  is the greatest angle.

So, the shortest side is BC.

**Answer4)** <u>Given</u>:  $\triangle ABC$ , side AB is produced to D so that BD = BC and  $\angle B = 60^{\circ}$ ,  $\angle A = 70^{\circ}$ 

#### To Prove:

(i) AD > CD

And, (ii) AD > AC

#### <u>Proof:</u>

First join C and D

Now,

In **ABC** 

 $\angle A + \angle B + \angle C = 180^{\circ}$ 

```
\angle C = 180^{\circ} - 70^{\circ} - 60^{\circ} = 50^{\circ}
```

 $\angle C = 50^{\circ}$   $\angle ACB = 50^{\circ}$  (i) And also in  $\triangle BDC$   $\angle DBC = 180^{\circ} - \angle ABC$   $= 180^{\circ} - 60^{\circ} = 120^{\circ}$ BD = BC (Given)

 $\angle BCD = \angle BDC$ 

Now,

 $\angle DBC + \angle BCD + \angle BDC = 180^{\circ}$ 

(Sum of all sides of triangle)

(Therefore, angle opposite to equal sides are equal)

(Therefore,  $\angle ABD$  is straight angle)

609

°0

(Sum of all angles of triangle)

 $120^{\circ} + \angle BCD + \angle BCD = 180^{\circ}$ 

 $2 \angle BCD = 180^{\circ} - 120^{\circ}$ 

 $2 \angle BCD = 60^{\circ}$ 

 $\angle BCD = 30^{\circ}$ 

Therefore,  $\angle BCD = \angle BDC = 30^{\circ}$  (ii)

Now, consider  $\triangle$ BDC,



 $\angle BAC = \angle DAC = 70^{\circ}$  (Given)

 $\angle BDC = \angle ADC = 30^{\circ} [From (ii)]$ 

 $\angle ACD = \angle ACB + \angle BCD$ 

 $= 50^{\circ} + 30^{\circ}$  [From (i) and (ii)]

 $= 80^{\circ}$ 

Now,

 $\angle ADC < \angle DAC < \angle ACD$ 

d s AC < DC < AD (Therefore, side opposite to greater angle is longer and smaller angle is smaller)

And,

AD > AC

Hence Proved.

We have,


 $\angle ACD > \angle DAC$ 

And,

 $\angle ACD > \angle ADC$ 

AD > DC

AD > AC (Therefore, side opposite to greater angle is longer and smaller angle is smaller)



# **TO PROVE:**

AD < BC

# PROOF:

 $\angle B < \angle A$ 

SO,

OA < OB ...(1) (SIDE OPPOSITE TO SMALLER ANGLE IS SMALL)

NOW,

∠C<∠D

SO,

...(2) (SIDE OPPOSITE TO SMALLER ANGLE IS SMALL) 0D < 0C NOW,

ADDING 1 AND 2

OA + OD < OB + OC

AD < BC

ADDING WE GET,

HENCE PROVED.


Answer6) Given:

In quadrilateral ABCD, AB smallest & CD is longest sides.

To Prove:∠A>∠C

&∠B>∠D

**Construction:** Join AC.



**CLASS IX** 

Mark the angles as shown in the figure..

#### **Proof:**

In  $\triangle ABC$ , AB is the shortest side.

BC > AB

∠2>∠4 ...(i)

[Angle opposite to longer side is greater]

In  $\triangle$ ADC, CD is the longest side

CD > AD

∠1>∠3 ...(ii)

[Angle opposite to longer side is greater]

Adding (i) and (ii), we have

 $\angle 2 + \angle 1 > \angle 4 + \angle 3$ 

⇒∠A>∠C

that Similarly, by joining BD, we can prove that

 $\angle B \ge \angle D$ 

Answer 7) To Prove: (AB + BC + CD + DA) > (AC + BD)

# **Proof:**


ABCD is a quad.Its diagonals are AC and BD.

In triangle ACB, AB + BC > AC...(1)

In triangle BDC, BC + CD > BD...(2)

In triangle ACD, AD + DC > AC...(3)

In triangle BAD, AB + AD > BD...(4)



#### **CLASS IX**

Adding 1,2,3 and 4,

AB + BC + BC + CD + AD + DC + AB + AD > AC + BD + AC + BD

2AB + 2BC + 2CD + 2AD > 2AC + 2BD

AB + BC + CD + AD > AC + BD.HENCE PROVED.

**Answer8)** Since, the sum of lengths of any two sides in a triangle should be greater than the length of third side

Therefore, In  $\triangle$  AOB, AB < OA + OB .....(i)

In  $\triangle$  BOC, BC < OB + OC .....(ii)

In  $\Delta$  COD, CD < OC + OD .....(iii)

In  $\triangle$  AOD, DA < OD + OA .....(iv)

Abooks, Mitch awa  $\Rightarrow$  AB + BC + CD + DA < 20A + 20B + 20C + 20D

 $\Rightarrow$  AB + BC + CD + DA < 2[(AO + OC) + (DO + OB)]

 $\Rightarrow$  AB + BC + CD + DA < 2(AC + BD)

Hence Proved.

**Answer9**) Given: In  $\triangle ABC$ ,  $\angle B = 35^{\circ}$ ,  $\angle C = 65^{\circ}$  and  $\angle BAX = \angle XAC$ 

To find: Relation between AX, BX and CX in descending order.

In  $\triangle$ ABC, by the angle sum property, we have


 $\angle A + \angle B + \angle C = 180^{\circ}$ 

 $\angle A + 35^{\circ} + 65^{\circ} = 180^{\circ}$ 

 $\angle A + 100^{\circ} = 180^{\circ}$ 

 $\therefore \angle A = 80^{\circ}$ 

But  $\angle BAX = \angle A = 40^{\circ}$ 



#### **CLASS IX**

# Vedantu

Now in  $\triangle ABX$ ,

 $\angle B = 35^{\circ}$ 

 $\angle BAX = 40^{\circ}$ 

And  $\angle BXA = 180^{\circ} - 35^{\circ} - 40^{\circ}$ 

= 105°

So, in  $\triangle ABX$ ,

 $\angle$ B is smallest, so the side opposite is smallest, i.e. AX is smallest side.

 $\therefore AX < BX \dots (1)$ 

Now consider  $\triangle AXC$ ,

 $\angle CAX = \angle A = 40^{\circ}$ 

 $\angle AXC = 180^{\circ} - 40^{\circ} - 65^{\circ}$ 

 $= 180^{\circ} - 105^{\circ} = 75^{\circ}$ 

Hence, in  $\triangle AXC$  we have,

 $\angle CAX = 40^\circ, \angle C = 65^\circ, \angle AXC = 75^\circ$ 

 $\therefore \angle CAX$  is smallest in  $\triangle AXC$ 


So the side opposite to  $\angle CAX$  is shortest

i.e. CX is shortest

 $\therefore$  CX <AX .... (2)


From 1 and 2,

BX > AX > CX



**Answer12)** Let  $\triangle$ ABC be a triangle in which AC is the longest side.

**To prove:** Angle opposite the longest side is greater than 2/3 of right angle.



AB + BC > AC ...(1)

In  $\triangle$ ADC, we have

CD + DA > AC ...(2)

Adding (1) and (2), we get

AB + BC + CD + DA > AC + AC

CD + DA + AB + BC > 2 AC

Hence Proved.

#### Answer14)

## Given:

In triangle ABC, O is any interior point.

We know that any segment from a point O inside a triangle to any vertex of the triangle cannot be longer than the two sides adjacent to the vertex.

Thus, OA cannot be longer than both AB and CA (if this is possible, then O is outside the triangle).

# To Prove:

- (i) AB + AC > OB + OC
- (ii) AB + BC + CA > OA + OB + OC

(iii) 
$$OA + OB + OC > \frac{1}{2} (AB + BC + CA)$$

# Proof:

(i) OA cannot be longer than both AB and CA AB>OB ...(1)

AC>0C ...(2)

Thus,

AB+AC>OB+OC ...[Adding (1) and(2)]

AB>0B ...(1)

# **CLASS IX**

# **RS Aggarwal solutions**

D

O

В

AC>0C ...(2)

Thus.

AB+AC>OB+OC ...[Adding (1) and(2)]

(ii) AB>0A.....(3)

BC>0B.....(4)

CA>0C.....(5) Adding the above three equations, we get:

Thus, AB+BC+CA>OA+OB+OC ...(6)

OA cannot be longer than both AB and CA.

د..., une first equation to be proved is shown correct. (iii) Now, consider the triangles OAC, OBA and OBC. We have: OA+OC>AC اA+OB>AB B+OC>BC

Adding the above three equations, we get:

OA+OC+OA+OB+OB+OC>AB+AC+BC

 $\Rightarrow$ 2(0A+0B+0C)>AB+AC+BC

Thus, OA+OB+OC>1/2(AB+BC+CA).

Answer15) Given : (i)  $AD \perp BC$ 

(ii) CD > BD

To Prove: AC > AB

# Proof:

In  $\triangle$  ABD ; $\angle$ ABD +  $\angle$ BAD +  $\angle$ BDA = 180°

 $\angle ABD + \angle BAD + 90^\circ = 180^\circ$ 

 $\angle ABD + \angle BAD = 90^{\circ}$ 

Similarly; In  $\triangle$ ADC ; $\angle$ ACB +  $\angle$ CAD = 90°

Since; BD <CD ;∠BAD <∠CAD

∠ABD >∠ACB

AC > AB

(sides opposite to greater angles are greater)

Answer16) <u>Given:</u> CD = DE

To prove: AB + AC > BE

Proof:

In  $\triangle ABC$ ,

AB+AC>BC ...(1)

AB+AC>BC ...1

In  $\triangle$ BED,

BD+CD>BE⇒BC>BE ...(2)



#### **RS** Aggarwal solutions

Α

D

C

R

BD+CD>BE⇒BC>BE ...2

From (1) and (2), we get

AB + AC > BE.

Hence Proved.


Same textbooks which and

# **MULTIPLE CHOICE QUESTIONS**

# Answer1) (a)

SSA is not a criteria for congruency of triangles. SSA would mean for example, that in triangles ABC and DEF, angle A = angle D, AB = DE, and BC = EF.

With these assumptions it is *not* true that triangle ABC is congruent to triangle DEF. In general there are two sets of congruent triangles with the same SSA data.




Ρ

Hence,  $\Delta PQR \cong \Delta CAB$ .


# Answer3 (a)

<u>**Given**</u> :  $\triangle ABC \cong \triangle PQR$ 

From the above given information following figures can be drawn.



From the above given information following figure can be drawn.




Since;  $\Delta$  ABC is an isosceles triangle Hence;  $\angle B = \angle C = 50^{\circ}$ 

 $\angle A + \angle B + \angle C = 180^{\circ}$  $\angle x + 50^{\circ} + 50^{\circ} = 180^{\circ}$ ∠x+100°=180° ∠x =180° - 100°  $\angle x = \angle A = 80^{\circ}$ 

Answer5) (a)

**<u>Given:</u>** (i) BC = AB(ii) ∠B =80°

POKS MISCH SMEN From the above given information following figure can be drawn.



Since;  $\Delta$  ABC is an isosceles triangle  $\angle A = \angle C = \angle x$ 

Hence;  $\angle A + \angle B + \angle C = 180^{\circ}$ 


$$\angle x + \angle x + 80^{\circ} = 180^{\circ}$$
$$2 \angle x = 180^{\circ} \cdot 80^{\circ}$$
$$\angle x = \frac{100^{\circ}}{2}$$
$$\angle x = 50^{\circ}$$
$$\angle x = \angle A = 50^{\circ}$$

#### Answer6) (a)

<u>Given:</u> (i)  $\angle A = \angle C$ 

- (ii) BC = 4 cm
- (iii) AC = 5 cm

From the above given information following figure can be drawn.



Hence;  $\Delta$  ABC is an isosceles triangle.

So; AB = BC = 4 cm.

#### Answer7) (b)

**<u>Given:</u>** (i) side 1 = 4 cm.

(ii) side 2 = 2.5 cm.

Since; the sum of two sides in a triangle must be greater than the third side.

So; the third side should be less than the sum of the other two sides.


the third side should be  $<4\ \mathrm{cm}+2.5\ \mathrm{cm}$  , i.e. 6.5 cm.

Hence; the third side cannot be 6.5 cm.


Answer 8) (b)


<u>Given</u> :  $\angle C > \angle B$ 

From the above given information following figure can be drawn.



From the above given information following figures can be drawn.





 $\angle A + \angle B + \angle C = 180^{\circ}$ 

 $40^{\circ} + 60^{\circ} + \angle C = 180^{\circ}$ 

 $\angle C = 180^{\circ} - 100^{\circ}$ 


$$\angle C = 80^{\circ}$$

Since; side opposite of greater angle is greater. So, the side AB is greatest.

Cestioon S. Hit

#### Answer11(c)

<u>Given:</u> AB > AC



D

So,  $AB > AC = \angle ACB > \angle ABC$ 

В

 $= \angle ACD > \angle ABD$ . ....(i)

Again side CD of  $\Delta$  ACD has been produced to B.



So, ext.  $\angle ADB > \angle ACD$ .

From (i) and (ii), we get

 $\angle ADB > \angle ACD > \angle ABD$ 

∠ADB >∠ABD

AB > AD (side opposite of greater angle is greater).

Hence, AB > AD.

# Answer12) (b)

<u>Given:</u> (i) AB > AC.

(ii) BO and CO are the bisectors of  $\angle B$  and  $\angle C$ .

 $\angle ACB > \angle ABC$ 

Angles opposite to greater sides are greater.

B

 $\frac{\angle ACB}{2} > \frac{\angle ABC}{2}$ 

 $\angle 0CB > \angle 0BC$ 

Hence, OB > OC.

Answer13) (a)

<u>Given:</u> (i) AB = AC

**CLASS IX** 


# **RS Aggarwal solutions**

Ο

HINCH BINDY

С

(ii) 
$$OB = OC$$




# Answer15) (b)

 $\underline{\text{Given}}$ : (i) AB = DE

(ii) BC = EF

From the above given information following figures can be drawn.



From the above given information following figures can be drawn.

For  $\triangle ABC \cong \triangle DEF$ , BC should be equal to EF. Hence if BC = EF, then  $\triangle ABC \cong \triangle DEF$  by A.S.A. criterion.

Answer17) (a) Given: (i) AB = AC(ii)  $\angle C = \angle P$ (iii)  $\angle B = \angle Q$ A A B = AC  $\angle C = \angle B$   $\angle P = \angle Q$  (since  $\angle C = \angle P$  and  $\angle B = \angle Q$ ) QR = PQ. Thus both the triangles are isosceles but not congruent.

# Answer18) (c)

Two right angles would up to 180° so the third angle becomes zero. This is not possible. Therefore, the triangle cannot have two right angles. A triangle can't have two obtuse angles as obtuse angle meAnswer more than 90°. So, the sum of the two sides exceeds more than 180° which is not possible. As the sum of all three angles of a triangle is 180°. A triangle can have two acute angle as acute angle meAnswer less than 90°. And *External angle* of *triangle* is greater *than either opposite angles*.

**Answer19)** a) (Sum of any sides of a triangle) **<u>greater than(>)</u>** (third side).

b) (Difference of any two sides of a triangle) **less than(<)** (third side).

c) (Sum of three altitudes of a triangle) less than(<) (sum of three sides).

- d) (Sum of any two sides of a triangle) **greater than(>)** (twice the median to the third side).
- e) (Perimeter of a triangle) **greater than(>)** (sum of its three median).

# **CLASS IX**

**Answer20)** a) Each angle of an equilateral triangle measures <u>60</u>°.

- b) MediAnswer of an equilateral triangle are equal.
- c) In a right angle triangle, the hypotenuse is the **longest** side.
- d) Drawing a  $\triangle$ ABC with AB = 3 cm., BC = 4 cm. and CA = 7 cm. is **not possible**.

