

**Exercise 12.2 : Solutions of Questions on Page Number : 273**

**Q1 :**

**Find the distance between the following pairs of points:**

(i) (2, 3, 5) and (4, 3, 1) (ii) (-3, 7, 2) and (2, 4, -1)

(iii) (-1, 3, -4) and (1, -3, 4) (iv) (2, -1, 3) and (-2, 1, 3)

**Answer :**

The distance between points  $P(x_1, y_1, z_1)$  and  $P(x_2, y_2, z_2)$  is given

by 
$$PQ = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$$

(i) Distance between points (2, 3, 5) and (4, 3, 1)

$$= \sqrt{(4-2)^2 + (3-3)^2 + (1-5)^2}$$


$$= \sqrt{(2)^2 + (0)^2 + (-4)^2}$$

$$= \sqrt{4+16}$$

$$= \sqrt{20}$$

$$= 2\sqrt{5}$$

(ii) Distance between points  $(-3, 7, 2)$  and  $(2, 4, -1)$



$$\begin{aligned}
&= \sqrt{(2+3)^2 + (4-7)^2 + (-1-2)^2} \\
&= \sqrt{(5)^2 + (-3)^2 + (-3)^2} \\
&= \sqrt{25+9+9} \\
&= \sqrt{43}
\end{aligned}$$

(iii) Distance between points  $(-1, 3)$  and  $(1, -3, 4)$

$$\begin{aligned}
&= \sqrt{(1+1)^2 + (-3-3)^2 + (4+4)^2} \\
&= \sqrt{(2)^2 + (-6)^2 + (8)^2} \\
&= \sqrt{4+36+64} = \sqrt{104} = 2\sqrt{26}
\end{aligned}$$

(iv) Distance between points  $(2, -1, 3)$  and  $(-2, 1, 3)$

$$\begin{aligned}
&= \sqrt{(-2-2)^2 + (1+1)^2 + (3-3)^2} \\
&= \sqrt{(-4)^2 + (2)^2 + (0)^2} \\
&= \sqrt{16+4} \\
&= \sqrt{20} \\
&= 2\sqrt{5}
\end{aligned}$$

**Q2 :**

**Show that the points  $(-2, 3, 5)$ ,  $(1, 2, 3)$  and  $(7, 0, -1)$  are collinear.**

**Answer :**

Let points  $(-2, 3, 5)$ ,  $(1, 2, 3)$ , and  $(7, 0, -1)$  be denoted by P, Q, and R respectively.

Points P, Q, and R are collinear if they lie on a line.

$$\begin{aligned}
 PQ &= \sqrt{(1+2)^2 + (2-3)^2 + (3-5)^2} \\
 &= \sqrt{(3)^2 + (-1)^2 + (-2)^2} \\
 &= \sqrt{9+1+4} \\
 &= \sqrt{14}
 \end{aligned}$$

$$\begin{aligned}
 QR &= \sqrt{(7-1)^2 + (0-2)^2 + (-1-3)^2} \\
 &= \sqrt{(6)^2 + (-2)^2 + (-4)^2} \\
 &= \sqrt{36+4+16} \\
 &= \sqrt{56} \\
 &= 2\sqrt{14}
 \end{aligned}$$

$$\begin{aligned}
 PR &= \sqrt{(7+2)^2 + (0-3)^2 + (-1-5)^2} \\
 &= \sqrt{(9)^2 + (-3)^2 + (-6)^2} \\
 &= \sqrt{81+9+36} \\
 &= \sqrt{126} \\
 &= 3\sqrt{14}
 \end{aligned}$$

Here,  $PQ + QR = \sqrt{14} + 2\sqrt{14} = 3\sqrt{14} = PR$

Hence, points P(â€“2, 3, 5), Q(1, 2, 3), and R(7, 0, â€“1) are collinear.

**Q3 :**

**Verify the following:**

- (i) (0, 7, -10), (1, 6, -6) and (4, 9, -6) are the vertices of an isosceles triangle.
- (ii) (0, 7, 10), (-1, 6, 6) and (-4, 9, 6) are the vertices of a right angled triangle.
- (iii) (-1, 2, 1), (1, -2, 5), (4, -7, 8) and (2, -3, 4) are the vertices of a parallelogram.

**Answer :**

- (i) Let points (0, 7, â€“10), (1, 6, â€“6), and (4, 9, â€“6) be denoted by A, B, and C respectively.

$$\begin{aligned}
 AB &= \sqrt{(1-0)^2 + (6-7)^2 + (-6+10)^2} \\
 &= \sqrt{(1)^2 + (-1)^2 + (4)^2} \\
 &= \sqrt{1+1+16} \\
 &= \sqrt{18} \\
 &= 3\sqrt{2}
 \end{aligned}$$

$$\begin{aligned}
 BC &= \sqrt{(4-1)^2 + (9-6)^2 + (-6+6)^2} \\
 &= \sqrt{(3)^2 + (3)^2} \\
 &= \sqrt{9+9} = \sqrt{18} = 3\sqrt{2}
 \end{aligned}$$

$$\begin{aligned}
 CA &= \sqrt{(0-4)^2 + (7-9)^2 + (-10+6)^2} \\
 &= \sqrt{(-4)^2 + (-2)^2 + (-4)^2} \\
 &= \sqrt{16+4+16} = \sqrt{36} = 6
 \end{aligned}$$

Here,  $AB = BC \neq CA$

Thus, the given points are the vertices of an isosceles triangle.

(ii) Let  $(0, 7, 10)$ ,  $(-1, 6, 6)$ , and  $(-4, 9, 6)$  be denoted by A, B, and C respectively.

$$\begin{aligned}
 AB &= \sqrt{(-1-0)^2 + (6-7)^2 + (6-10)^2} \\
 &= \sqrt{(-1)^2 + (-1)^2 + (-4)^2} \\
 &= \sqrt{1+1+16} = \sqrt{18} \\
 &= 3\sqrt{2}
 \end{aligned}$$

$$\begin{aligned}
 BC &= \sqrt{(-4+1)^2 + (9-6)^2 + (6-6)^2} \\
 &= \sqrt{(-3)^2 + (3)^2 + (0)^2} \\
 &= \sqrt{9+9} = \sqrt{18} \\
 &= 3\sqrt{2}
 \end{aligned}$$

$$\begin{aligned}
 CA &= \sqrt{(0+4)^2 + (7-9)^2 + (10-6)^2} \\
 &= \sqrt{(4)^2 + (-2)^2 + (4)^2} \\
 &= \sqrt{16+4+16} \\
 &= \sqrt{36} \\
 &= 6
 \end{aligned}$$

$$\text{Now, } AB^2 + BC^2 = (3\sqrt{2})^2 + (3\sqrt{2})^2 = 18+18 = 36 = AC^2$$

Therefore, by Pythagoras theorem, ABC is a right triangle.

Hence, the given points are the vertices of a right-angled triangle.

(iii) Let  $(\hat{a}\hat{e}^{-1}, 2, 1)$ ,  $(1, \hat{a}\hat{e}^{-2}, 5)$ ,  $(4, \hat{a}\hat{e}^{-7}, 8)$ , and  $(2, \hat{a}\hat{e}^{-3}, 4)$  be denoted by A, B, C, and D respectively.

$$\begin{aligned}
 AB &= \sqrt{(1+1)^2 + (-2-2)^2 + (5-1)^2} \\
 &= \sqrt{4+16+16} \\
 &= \sqrt{36} \\
 &= 6
 \end{aligned}$$

$$\begin{aligned}
 BC &= \sqrt{(4-1)^2 + (-7+2)^2 + (8-5)^2} \\
 &= \sqrt{9+25+9} = \sqrt{43}
 \end{aligned}$$

$$\begin{aligned}
 CD &= \sqrt{(2-4)^2 + (-3+7)^2 + (4-8)^2} \\
 &= \sqrt{4+16+16} \\
 &= \sqrt{36} \\
 &= 6
 \end{aligned}$$

$$\begin{aligned}
 DA &= \sqrt{(-1-2)^2 + (2+3)^2 + (1-4)^2} \\
 &= \sqrt{9+25+9} = \sqrt{43}
 \end{aligned}$$

Here,  $AB = CD = 6$ ,  $BC = AD = \sqrt{43}$

Hence, the opposite sides of quadrilateral ABCD, whose vertices are taken in order, are equal.

Therefore, ABCD is a parallelogram.

Hence, the given points are the vertices of a parallelogram.

**Q4 :**

Find the equation of the set of points which are equidistant from the points (1, 2, 3) and (3, 2, -1).

**Answer :**

Let P (x, y, z) be the point that is equidistant from points A(1, 2, 3) and B(3, 2, -1).

Accordingly, PA = PB

$$\Rightarrow PA^2 = PB^2$$

$$\Rightarrow (x-1)^2 + (y-2)^2 + (z-3)^2 = (x-3)^2 + (y-2)^2 + (z+1)^2$$

$$\Rightarrow x^2 - 2x + 1 + y^2 - 4y + 4 + z^2 - 6z + 9 = x^2 - 6x + 9 + y^2 - 4y + 4 + z^2 + 2z + 1$$

$$\Rightarrow -2x - 4y - 6z + 14 = -6x - 4y + 2z + 14$$

$$\Rightarrow -2x - 6z + 6x - 2z = 0$$

$$\Rightarrow 4x - 8z = 0$$

$$\Rightarrow x - 2z = 0$$

Thus, the required equation is  $x - 2z = 0$ .

**Q5 :**

Find the equation of the set of points P, the sum of whose distances from A (4, 0, 0) and B (-4, 0, 0) is equal to 10.

**Answer :**

Let the coordinates of P be (x, y, z).

The coordinates of points A and B are (4, 0, 0) and (-4, 0, 0) respectively.

It is given that PA + PB = 10.

$$\Rightarrow \sqrt{(x-4)^2 + y^2 + z^2} + \sqrt{(x+4)^2 + y^2 + z^2} = 10$$

$$\Rightarrow \sqrt{(x-4)^2 + y^2 + z^2} = 10 - \sqrt{(x+4)^2 + y^2 + z^2}$$

On squaring both sides, we obtain

$$\Rightarrow (x-4)^2 + y^2 + z^2 = 100 - 20\sqrt{(x+4)^2 + y^2 + z^2} + (x+4)^2 + y^2 + z^2$$

$$\Rightarrow x^2 - 8x + 16 + y^2 + z^2 = 100 - 20\sqrt{x^2 + 8x + 16 + y^2 + z^2} + x^2 + 8x + 16 + y^2 + z^2$$

$$\Rightarrow 20\sqrt{x^2 + 8x + 16 + y^2 + z^2} = 100 + 16x$$

$$\Rightarrow 5\sqrt{x^2 + 8x + 16 + y^2 + z^2} = (25 + 4x)$$

On squaring both sides again, we obtain

$$25(x^2 + 8x + 16 + y^2 + z^2) = 625 + 16x^2 + 200x$$

$$\Rightarrow 25x^2 + 200x + 400 + 25y^2 + 25z^2 = 625 + 16x^2 + 200x$$

$$\Rightarrow 9x^2 + 25y^2 + 25z^2 - 225 = 0$$

Thus, the required equation is  $9x^2 + 25y^2 + 25z^2 - 225 = 0$ .

